


FORWARD HADRONIC SCATTERING FROM FEW GeV TO MULTI TeV WITHIN REGGE THEORY

J.R. Peláez

Departamento de Física Teórica II. Universidad Complutense. 28040 Madrid. Spain.



The Regge description of forward hadronic scattering is extended from ~ 1 GeV above each threshold up to the multi TeV range. This is achieved with a simple parametrization, that includes mass effects, a logarithmic growth based on an improved unitarity bound at intermediate energies and a separate factorization of singularities. This parametrization can be easily implemented for phenomenological use but also sheds light on subdominant trajectories and the Pomeron logarithmic growth law.

1 Introduction

In recent works¹ we have shown how it was possible to extend down to roughly 1 GeV above threshold the Regge description of those combinations of total cross sections that involved the Pomeron, P' and ρ Regge trajectories only. In particular we provided several fits to the $(\bar{p}p+pp)$, (K^+p+K^-p) , $\pi^\pm N$ and $\pi\pi$ cross sections.

I report here on *preliminary results*² that include the a and ω trajectories and extend the analysis to the total hadronic cross sections of $\bar{p}p$, pp , $\bar{p}n$, pn , $K^\pm p$, $K^\pm n$, $\pi^\pm N$ and $\pi\pi$, as well as the ImF/ReF ratios of $\bar{p}p$, pp , pn , $\pi^\pm N$ and $K^\pm p$ forward elastic amplitudes, F . The data has been obtained from the extensive compilation of the COMPAS group. Given the fact that the original references did not treat the systematic uncertainties uniformly, we have followed two fitting strategies: First, we keep the original uncertainties as such. This allows for an easy comparison with previous works, including the PDG and the extensive ones of the COMPETE group³. However, this introduces a bias toward those data sets, usually the oldest, that do not provide systematic uncertainties. Indeed, many of these data are incompatible within their statistical errors, and cannot be described simultaneously. This produces an artificially large $\chi^2/d.o.f.$ no matter what function is fitted. For that reason, in our second strategy, we have added a systematic error, but only to those data without it, of 0.5% for pp , 1% for $\bar{p}p$ and 1.5%

for other processes. The size of these additional errors has been chosen of the same order of magnitude given by other experiments for the same process, so that we give similar weight to all sets without discarding any point. To account for different ways of combining statistical and systematic errors, in the first strategy we have added them in quadrature and linearly in the second. In addition, we use σ^{total} data¹⁰ on $\pi^+\pi^-$, $\pi^-\pi^-$, $\pi^+\pi^0$, above 1.42GeV, plus one data point per channel reconstructed from phase shift analysis¹ at 1.42GeV. If using $\pi\pi$ low energy information^{1,2}, the ρ residue and intercept come out somewhat smaller and larger, respectively.

The contributions to the $F_{AB \rightarrow AB}$ amplitudes from the Pomeron P , the f (or P'), ρ , ω and a trajectories, are

$$\begin{aligned} F_{p^\pm p} &= (P_{NN} + f_{NN} + a_{NN} \mp \omega_{NN} \mp \rho_{NN})/2, & F_{p^\pm n} &= (P_{NN} + f_{NN} - a_{NN} \mp \omega_{NN} \pm \rho_{NN})/2, \\ F_{K^\pm p} &= (P_{KN} + f_{KN} + a_{KN} \mp \omega_{KN} \mp \rho_{KN})/2, & F_{K^\pm n} &= (P_{KN} + f_{KN} - a_{KN} \mp \omega_{KN} \pm \rho_{KN})/2, \\ F_{\pi^\pm p} &= (P_{\pi N} + f_{\pi N})/\sqrt{6} \mp \rho_{\pi N}/2, & F_{\pi^\pm \pi^-} &= (P_{\pi\pi} + f_{\pi\pi})/3 \pm \rho_{\pi\pi}/2, & F_{\pi^0 \pi^-} &= (P_{\pi\pi} + f_{\pi\pi})/3, \end{aligned}$$

where $N = p^\pm, n$ and isospin Clebsch Gordan coefficients have been extracted to simplify the factorization⁸ relations $R_{AB}(\nu) = f_A^R f_B^R R(\nu)$, where for $R = \rho, f, a, \omega$

$$R(\nu) = \beta_R \left(\frac{1 + \tau e^{-i\pi\alpha}}{\sin \pi\alpha} \right) \nu^{\alpha_R}, \quad (1)$$

where τ is the signature of the trajectory. If we want to take account of the masses, *the energy dependence should appear through* $\nu = (s - u)/2$, instead of the usual choice of s . Note that for forward scattering $\nu = s - m_a^2 - m_b^2 > s - s_{th}$. For the intercepts α_R we will assume, following the QCD version of Regge theory, and the recent analysis by the COMPETE group³, that the f/a and ρ/ω trajectories are degenerate, that is, that $\alpha_a = \alpha_f$ and $\alpha_\omega = \alpha_\rho$. In addition, some factors are redundant and can be absorbed by setting $f_\pi^R = 1$, for $R = P, f, \rho$ and $\beta_R = 1$ for $R = a, \omega$. This choice eases the notation and the comparison with previous works¹.

For the Pomeron, we propose the use of a “constant plus logarithm”, i.e.,

$$P_{AB} = C_{AB} + L_{AB}, \quad \text{Im } P(\nu) = \nu \left(\beta_P + A \log^2 \left[\frac{\nu - \nu_{th}}{\nu_1 \log^{7/2}(\nu/\nu_2)} \right] \right), \quad (2)$$

where the logarithmic law follows an improved unitarity bound⁹. The choice $\nu - \nu_{th}$ yields slightly better fits. Note that ν_{th} is ν at the branch point of the right cut for each amplitude. Indeed, it has been confirmed³ that the σ^{tot} growth and the $\text{Re } F/\text{Im } F$ ratios, are better described with $s \log s$ or, slightly better³, with $s \log^2 s$ terms rather than with $\alpha_P > 1$. Our Eq.(2) grows faster than $s \log s$ but slower at intermediate energies than the $s \log^2 s$ Froissart bound, which is nevertheless recovered at very high s . The recent generalization of the “factorization theorem”⁶ requires each singularity to factorize separately. Thus, as a first approximation, and for the Pomeron, we use separated factors: f_A^C and f_A^L .

Finally, the real parts of amplitudes are obtained from the dispersive representation, whereas total cross sections are given by:

$$\sigma_{ab} = 4\pi^2 \text{Im } F_{a+b \rightarrow a+b}(s, 0) / \lambda^{1/2}(s, m_a^2, m_b^2), \quad \lambda(s, m_a^2, m_b^2) = s^2 + (m_a^2 - m_b^2)^2 - 2s(m_a^2 + m_b^2)$$

However, λ is usually approximated by s^2 . Only very recently⁷ a slight improvement in χ^2/dof has been found by keeping the whole λ , instead of s^2 , down to $\sqrt{s} = 5$ GeV. Let us remark that, if $E_{kin} \simeq 1$ GeV, the effect of using s^2 , instead of λ , yields a 30% overestimation for NN .

2 Results

We show in Table 1 the parameters of the fits to data for different strategies and different E_{kin}^{min} , including the nominal uncertainty obtained from the χ^2/dof minimization routine MINUIT. Since the parameters are very strongly correlated these errors should be taken only nominally around the central values provided for strategy 2. Since there are flat directions in parameter space it is possible to get somewhat different parameter sets with almost the same χ^2/dof but beyond those nominal errors. An indication of the systematic error *in the determination of single parameters* can be obtained from the difference between strategies.

E_{kin}^{min}	strategy 2 1-1.3 GeV	strategy 1 1-1.3 GeV	Minuit errors	E_{kin}^{min}	strategy 2 1-1.3 GeV	strategy 1 1-1.3 GeV	Minuit errors
β_P	0.746	0.937	0.003	f_K^f	0.30	0.32	0.01
f_N^P	1.792	1.705	0.007	α_f	0.646	0.640	0.002
f_K^P	0.731	0.714	0.004	f_N^a	-0.24	0.25	0.04
A	0.043	0.050	0.001	f_K^a	-0.55	0.5	0.1
ν_1	0.0005	0.001	0.0001	β_ρ	1.28	1.34	0.11
ν_2	0.676	0.633	0.001	f_N^ρ	0.51	0.46	0.04
f_N^{log}	1.02	0.993	0.001	f_K^ρ	0.49	0.54	0.04
f_K^{log}	0.723	0.733	0.012	α_ρ	0.464	0.464	0.003
β_f	1.70	1.77	0.014	f_N^ω	1.97	1.98	0.015
f_N^f	1.78	1.75	0.01	f_K^ω	0.66	0.65	0.01
				σ_{LHC}	109 mb	110 mb	1mb

Table 1. Fit parameters with different strategies. The Minuit errors are *just statistical*, and nominal, since the parameters are strongly correlated and can only be used with the central values of *strategy 2*.

E_{kin}^{min} (GeV)	1-1.3	1.5	2	3
# data points	1186	1002	895	768
Parametrization	$\chi^2/d.o.f.$ for strategy 2 / 1			
Ours	0.85/1.56	0.63/1.14	0.57/1.05	0.52/0.95
$\nu_1 \equiv 0.01 \text{ GeV}^2$	0.85/1.57	0.63/1.26	0.58/1.06	0.52/0.97
powers of s^α	1.58/2.87	1.16/2.11	0.99/1.80	0.78/1.42
Pomeron logarithmic term				
$\nu \log(\nu)$	1.01/1.83	0.69/1.26	0.59/1.09	0.52/0.97
$\nu \log(\nu - \nu_{th})$	1.03/1.83	0.69/1.27	0.59/1.12	0.52/0.98
$\nu \log^2(\nu)$	0.97/1.79	0.68/1.24	0.59/1.10	0.52/0.95
$\nu \log^2(\nu - \nu_{th})$	0.91/1.68	0.65/1.18	0.58/1.06	0.52/0.95
Factorization of Pomeron logarithms				
$f_a^L \equiv 1$ (as PDG)	0.92/1.70	0.66/1.23	0.59/1.10	0.54/1.01
$f_a^C = f_a^L$	0.89/1.67	0.64/1.41	0.60/1.14	0.58/1.02

Table 2. $\chi^2/d.o.f.$ for several E_{kin}^{min} and different modifications of our parametrization.

In Table 2 we compare the $\chi^2/d.o.f.$ of parameterizations where we have not implemented some of our suggestions. Let us recall that these are: i) using ν instead of s , ii) the Eq.(2), iii) the separate factorization, i) and the correct flux factor. For each one of them *we find an improvement in $\chi^2/d.o.f.$* , with both fitting strategies.

In Figure 1 we show the curves obtained from our parametrization including, as gray bands, the nominal uncertainties from strategy 2. The dashed lines correspond to the PDG2004 parametrization⁴, valid above $\sqrt{s} = 5 \text{ GeV}$, but naively extrapolated below. We see that the simple parametrization reported here describes remarkably well 20 observables extending

from several TeV down to ~ 1 GeV above the threshold of each reaction. Further details will be given in a forthcoming publication², but apart from establishing the logarithmic growth of the Pomeron, we hope it could be easily used for dispersive studies in hadronic physics that involve integrals from the resonance region to infinity.




Figure 1: Results from our fit down to 1 GeV above threshold. Total NN , $\pi\pi$, $\pi^\pm N$, $K^\pm p$ and $K^\pm n$ cross sections as a function of $\sqrt{s}$, with detail of pp and $\bar{p}p$ at low energies. In the bottom row we show the results for $Re F/Im F$. The bands cover the nominal uncertainties in the parameters.

References

1. J. R. Pelaez and F. J. Yndurain, Phys. Rev. D **68**, 074005 (2003); Phys. Rev. D **69**, 114001 (2004) and hep-ph/0411334.
2. J.R. Pelaez, in preparation.
3. J. R. Cudell *et al.*, (COMPETE Collab.) Phys. Rev. D **65**, 074024 (2002)
4. *The review of Particle Physics* S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004). K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002). D.E. Groom *et al.*, Eur. Phys. J. C **15** (2000) 1.
5. Froissart, M. (1961). Proc. la Jolla Conf. on Elementary Particles; Gribov, V. N. (1962). Sov. Phys. JETP **14**, 1395.
6. J. R. Cudell, E. Martynov and G. Soyez, Nucl. Phys. B **682**, 391 (2004)
7. J. R. Cudell, E. Martynov, O. Selyugin and A. Lengyel, Phys. Lett. B **587**, 78 (2004)
8. Gell-Mann, M. Phys. Rev. Lett., **8**, 263, (1962); Gribov, V. N., and Pomeranchuk, I. Ya. Phys. Rev. Lett., **8**, 343, (1962).
9. F. J. Yndurain, Phys. Lett. B **41**, 591 (1972). (In that paper $\log^7$ was incorrectly given instead of $\log^{7/2}$).

10. Biswas, N. N., et al., *Phys. Rev. Lett.*, **18**, 273 (1967) Cohen, D. et al., *Phys. Rev.* **D7**, 661 (1973). Robertson, W. J., Walker, W. D., and Davis, J. L., *Phys. Rev.* **D7**, 2554 (1973). Hanlon, J., et al, *Phys. Rev. Lett.*, **37**, 967 (1976); Abramowicz, H., et al. *Nucl. Phys.*, **B166**, 62 (1980)